

Negative Magneto-Resistance Beyond Weak Localization in Three-Dimensional Billiards: Effect of Arnold Diffusion

Jun Ma and Katsuhiro Nakamura

Department of Applied Physics, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan

(February 1, 2008)

Abstract

We investigate a semiclassical conductance for ballistic open three-dimensional (3-d) billiards. For partially or completely broken-ergodic 3-d billiards such as $SO(2)$ symmetric billiards, the dependence of the conductance on the Fermi wavenumber is dramatically changed by the lead orientation. Application of a symmetry-breaking weak magnetic field brings about mixed phase-space structures of 3-d billiards which ensures a novel Arnold diffusion that cannot be seen in 2-d billiards. In contrast to the 2-d case, the anomalous increment of the conductance should inevitably include a contribution arising from Arnold diffusion as well as a weak localization correction. Discussions are devoted to the physical condition for observing this phenomenon.

PACS numbers: 03.65Sq; 05.45.+b; 73.20.Dx.

I. INTRODUCTION

Concave and convex billiards as prototypes of conservative chaotic systems have received a growing theoretical and experimental interest in the fields of nonlinear dynamics and statistical mechanics [1]. These billiards are nowadays fabricated as quantum dots or antidots in ballistic microstructures where the system size is much less than the mean free path ℓ ($\sim 20\mu\text{m}$) and larger than the Fermi wavelength ($\lambda \sim 50\text{nm}$)[2]. Since the striking experiments on the magneto-conductance of nanoscale stadium and circle billiards with a pair of conducting leads [2], quantum transport phenomena have been intensively investigated [3]. Some salient features of experiments well agreed with the semiclassical theory, where a semiclassical conductance formula was derived for two-dimensional (2-d) chaotic billiards by combining the semiclassical Green function with the Landauer formula [4,5]. Subsequently, the semiclassical conductance formula was extended for integrable 2-d billiards [6], and detailed numerical computation on circle billiards indicated that the conductance also depends on the lead orientation [7]. Recently, the dependence of the conductance on the lead orientation was found experimentally in a ballistic square billiard [8].

At present, another important object of research on quantum transport is the three-dimensional (3-d) billiard [9]. The ballistic 3-d quantum dots will be fabricated, for instance, by exploiting drying etching processes with focused-ion beams applied to $\text{Al}_x\text{Ga}_{1-x}\text{As}/\text{GaAs}/\text{Al}_x\text{Ga}_{1-x}\text{As}$ double hetero-structures, which will be less than both the elastic mean free path and phase coherence length. The subject of this paper is to extend the semiclassical conductance formula for 2-d billiards to 3-d billiards and to investigate the quantum transport in open 3-d billiards from a viewpoint of the Arnold diffusion, a key concept in high-dimensional nonlinear dynamical systems. While Arnold diffusion is a classical phenomenon, an anomalous phenomenon like this in classical mechanics should have a quantum counterpart, which can only be analytically revealed by using the semiclassical theory.

II. SEMICLASSICAL CONDUCTANCE FOR OPEN 3-DIMENSIONAL BILLIARDS

The conductance for open 3-d billiards connected to a pair of rectangular parallel-piped lead wires can be expressed by the transmission coefficient T as $G = \frac{2e^2}{h}T = \frac{2e^2}{h} \sum_{\mathbf{n}, \mathbf{m}=1} |t_{\mathbf{n}\mathbf{m}}|^2$, where $t_{\mathbf{n}\mathbf{m}}$ is the $\mathbf{S}$ -matrix element between the incoming mode $\mathbf{m} = (m_x, m_y)$ and the outgoing mode $\mathbf{n} = (n_x, n_y)$, and the double summation is taken over all propagating modes. The transmission amplitude connecting those modes at the Fermi energy E_F reads as [10,11]

$$t_{\mathbf{n}\mathbf{m}} = -i\hbar\sqrt{v_{\mathbf{n}}v_{\mathbf{m}}}\int dx''dy''dx'dy'\psi_{\mathbf{n}}^*(x'',y'')G(x'',y'',z'';x',y',z';E_F)\psi_{\mathbf{m}}(x',y'), \quad (1)$$

where $v_{\mathbf{m}}$ and $v_{\mathbf{n}}$ are the longitudinal velocities of electrons for the incoming and outgoing modes, respectively. x', y', z' and x'', y'', z'' are the *local* coordinates for the transverse (x, y) and longitudinal (z) directions (inward the billiard for z' and outward for z'') of the incoming and outgoing leads, respectively, and $G(x'', y'', z''; x', y', z'; E_F)$ is the Green function for an electron propagating from the entrance to the exit. $\psi_{\mathbf{m}}(x', y')$ and $\psi_{\mathbf{n}}(x'', y'')$ are the transverse components of the wave functions at the leads. To simplify the problem, the cross section of the leads is assumed to be a square with side length l . Then $\psi_{\mathbf{m}}(x', y') = \frac{2}{l} \sin(m_x \pi x'/l) \sin(m_y \pi y'/l)$ and $\psi_{\mathbf{n}}(x'', y'') = \frac{2}{l} \sin(n_x \pi x''/l) \sin(n_y \pi y''/l)$. In the ballistic regime, the Green function from entrance ($\mathbf{r}'$) to exit ($\mathbf{r}''$) can be replaced by the Gutzwiller semiclassical path-integral expression [12]

$$G(x'', y'', z''; x', y', z'; E_F) = \frac{2\pi}{(2\pi i\hbar)^2 v_F} \sum_{s(\mathbf{r}', \mathbf{r}'')} |\det(\frac{\partial \mathbf{p}'_{\perp}}{\partial \mathbf{r}''_{\perp}})|^{1/2} \exp(\frac{i}{\hbar} S_s(\mathbf{r}'', \mathbf{r}', E_F) - i\pi\nu_s/2), \quad (2)$$

where S_s and ν_s are the action and Maslov index for a trajectory s , and v_F is the Fermi velocity. The above semiclassical Green function was widely applied to derive the level density of both chaotic and regular billiards [12,13]. Here, to avoid direct transmission without any bouncing at the billiard wall we may place a stopper inside the billiard.

In the integration in Eq.(1) we introduce the entrance angle θ' between the initial direction of the orbit and the z' axis, and the angle φ' between the projection of the initial direction onto the $x'-y'$ plane and the x' axis. Angles θ'' and φ'' are defined for the exit in the same way. Then, we can rewrite $|\det(\frac{\partial \mathbf{p}'}{\partial \mathbf{r}_\perp})|$ for each trajectory s as $|\det(\frac{\partial p_\perp^{1'}, p_\perp^{2'}}{\partial (r_\perp^{1'}, r_\perp^{2'})})| = p_F^2 A_s$ where $A_s = |\frac{\sin \theta'}{\cos \theta''} (\frac{\partial \varphi'}{\partial x''} \frac{\partial \theta'}{\partial y''} - \frac{\partial \varphi'}{\partial y''} \frac{\partial \theta'}{\partial x''})|$. For large mode numbers, the integration over the transverse coordinates in Eq.(1) can be performed in the stationary-phase approximation, and the transmission amplitude becomes

$$t_{\mathbf{nm}} = -i\hbar \frac{16\pi}{l^2 p_F} \sum_{s(\bar{m}_x, \bar{m}_y, \bar{n}_x, \bar{n}_y)} \text{sgn}(\bar{m}_x) \text{sgn}(\bar{m}_y) \text{sgn}(\bar{n}_x) \text{sgn}(\bar{n}_y) (\cos \theta' \cos \theta'')^{1/2} (A_s/B_s)^{1/2} e^{\frac{i}{\hbar} \tilde{S}_s - i\tilde{\nu}_s \pi/2} \quad (3)$$

with

$$B_s = [|\sin 2\varphi'' (\frac{\partial \theta''}{\partial y''} \frac{\partial \theta''}{\partial x''} \cos^2 \theta'' - \frac{\partial \varphi''}{\partial y''} \frac{\partial \varphi''}{\partial x''} \sin^2 \theta'') + \sin 2\theta'' (\frac{\partial \varphi''}{\partial y''} \frac{\partial \theta''}{\partial x''} \cos^2 \varphi'' - \frac{\partial \theta''}{\partial y''} \frac{\partial \varphi''}{\partial x''} \sin^2 \varphi'')| \\ \times |\sin 2\varphi' (\frac{\partial \theta'}{\partial y'} \frac{\partial \theta'}{\partial x'} \cos^2 \theta' - \frac{\partial \varphi'}{\partial y'} \frac{\partial \varphi'}{\partial x'} \sin^2 \theta') + \sin 2\theta' (\frac{\partial \varphi'}{\partial y'} \frac{\partial \theta'}{\partial x'} \cos^2 \varphi' - \frac{\partial \theta'}{\partial y'} \frac{\partial \varphi'}{\partial x'} \sin^2 \varphi')|], \quad (4)$$

where $\bar{m}_x = \pm m_x, \bar{m}_y = \pm m_y, \bar{n}_x = \pm n_x, \bar{n}_y = \pm n_y$, and $\tilde{S}_s = S(\mathbf{r}''_s, \mathbf{r}'_s, E_F) + \hbar\pi\bar{m}_x x'_s/l + \hbar\pi\bar{m}_y y'_s/l - \hbar\pi\bar{n}_x x''_s/l - \hbar\pi\bar{n}_y y''_s/l$, and $\tilde{\nu}$ is the Maslov index which includes an extra phase coming from the possible sign change in each of the four Fresnel integrals. The above result means that only those isolated trajectories connecting a pair of transverse planes at discrete angles $\sin \theta' \cos \varphi' = \bar{m}_x \pi/k_F l, \sin \theta' \sin \varphi' = \bar{m}_y \pi/k_F l, \sin \theta'' \cos \varphi'' = \bar{n}_x \pi/k_F l$, and $\sin \theta'' \sin \varphi'' = \bar{n}_y \pi/k_F l$ dominate the conductance.

Using Eqs.(3,4), we shall first analyze a completely chaotic 3-d billiard with a pair of conducting leads. In this case, the electron injected from the incoming lead will bounce at the billiard boundary in an ergodic way before reaching the exit, leading to amplitudes of the same order for both transmission and reflection coefficients. In the large mode number case, the summations over modes are replaced by integrations and the transmission coefficient $T = \sum_{\mathbf{n}, \mathbf{m}} |t_{\mathbf{nm}}|^2$ is rewritten as

$$T = \hbar^2 \frac{256\pi^2}{l^4} \frac{1}{p_F^2} \sum_{\bar{m}_x, \bar{m}_y, \bar{n}_x, \bar{n}_y} \sum_{s, u} \cos \theta' \cos \theta'' (\frac{A_s A_u}{B_s B_u})^{1/2} e^{\frac{i}{\hbar} (\tilde{S}_s - \tilde{S}_u) - i\frac{\pi}{2} (\tilde{\nu}_s - \tilde{\nu}_u)}$$

$$\begin{aligned}
& \approx \frac{256\pi^2}{l^4 k_F^2} \int_0^{\sqrt{\bar{n}_x^2 + \bar{n}_y^2} \leq \frac{k_F^2 l^2}{\pi^2}} d\bar{n}_x d\bar{n}_y \int_0^{\sqrt{\bar{m}_x^2 + \bar{m}_y^2} \leq \frac{k_F^2 l^2}{\pi^2}} d\bar{m}_x d\bar{m}_y \sum_{s,u} \cos \theta' \cos \theta'' \\
& \times \left(\frac{A_s A_u}{B_s B_u} \right)^{\frac{1}{2}} e^{\frac{i}{\hbar}(\tilde{S}_s - \tilde{S}_u) - i\frac{\pi}{2}(\tilde{\nu}_s - \tilde{\nu}_u)} \\
& = \frac{256k_F^2}{\pi^2} \int_{-1}^1 \sin \theta' d(\sin \theta') \int_{-\pi}^{\pi} d\varphi' \int_{-1}^1 \sin \theta'' d(\sin \theta'') \int_{-\pi}^{\pi} d\varphi'' \sum_{s,u} \cos \theta' \cos \theta'' \\
& \times \left(\frac{A_s A_u}{B_s B_u} \right)^{\frac{1}{2}} e^{\frac{i}{\hbar}(\tilde{S}_s - \tilde{S}_u) - i\frac{\pi}{2}(\tilde{\nu}_s - \tilde{\nu}_u)}. \tag{5}
\end{aligned}$$

Noting the dimension of $(\frac{A_s A_u}{B_s B_u})^{1/2}$ to be l^2 , and using the diagonal approximation, the above integration can be evaluated as $T \propto (k_F l)^2$. The conductance for 3-d chaotic billiards is thus given by $G \propto \frac{e^2}{h} (k_F l)^2$ without any dependence on the lead orientations.

III. COMPLETELY OR PARTIALLY BROKEN-ERGODIC 3-D BILLIARDS

However, the situation will be dramatically changed for the case of completely or partially broken-ergodic 3-d billiards. As examples, we choose SO(2)-symmetric billiards: A completely integrable (: broken-ergodic) 3-d billiard is available by rotating, e.g., the 2-d ellipse billiard (on the $Y - Z$ plane) around the Z axis (see Fig.1). Note: X, Y, Z are the *global* coordinates; A partially broken-ergodic 3-d billiard is also available in a similar way by rotating, e.g., the 2-d stadium. All these SO(2)-symmetric 3-d billiards have the angular momentum L_Z as a constant of motion. In an open system version of each of these billiards, one may place in its inside a smaller stopper of the similar shape, to avoid direct transmission of electrons. The resultant 3-d *shell* billiards still retain the SO(2) symmetry.

Firstly, consider the vertical case (i) when the incoming and outgoing leads are connected with the billiard vertically at points A and C , respectively (see Fig.1). Electrons to reach the exit C should have the vanishing angular momentum, $L_Z = 0$. Therefore, only the electrons with the initial velocity vector at the entrance A lying in the $Y - Z$ plane can reach C because of the conservation of L_Z ; Other incoming electrons falling into the trajectories out of this plane cannot reach C and should return to A . In deriving the transmission coefficient for this lead orientation, the integration prior to Eq.(5) is reduced to the line integration

A 3D diagram showing a horizontal elliptical disk in a Cartesian coordinate system with axes X , Y , and Z . The Z -axis is vertical, and the Y -axis is horizontal to the right. A curved arrow around the Z -axis indicates rotation. A vertical line with two parallel arrows pointing downwards is labeled c at the top and Q at the bottom. On the left edge of the disk, a point is labeled A , and a horizontal line segment is labeled I . Three arrows point away from the bottom of the disk towards the label B .

over θ' and θ'' only, yielding the conductance $G \propto \frac{e^2}{h} k_F l$ (linear in $k_F l$). (The reflection coefficient, which is obtained by applying the same procedure as in the chaotic billiard, is proportional to $(k_F l)^2$.) We shall then consider the parallel case (ii) when the incoming and outgoing leads are parallelly placed at points Q and C , respectively (see Fig.1). Each electron incoming from the entrance Q has the vanishing angular momentum, $L_Z = 0$, and thereby can reach C or return to Q with probability of the same order. Therefore, for this lead orientation, $G \propto \frac{e^2}{h} (k_F l)^2$.

6

by the lead orientation. It should be noted: open 2-d billiards always share $G \propto \frac{e^2}{h} k_F l$ [4], independently of both integrability and lead orientations.

IV. EFFECTS OF SYMMETRY-BREAKING WEAK MAGNETIC FIELD

The magneto-resistance is also a very important problem in 3-d billiards. It is well known that in ballistic 2-d billiards the decrease of resistance with increasing magnetic field (: negative magneto-resistance) occurs due to the weak localization based on the quantum interference between a pair of time-reversal symmetric orbits[2-5]. What kind of additional novel phenomena will be expected in SO(2)-symmetric 3-d billiards with the vertical lead (A, C) orientation when a symmetry-breaking weak magnetic field B will be applied along the X axis? In contrast to phase-space structures in 2-d systems where each of chaotic zones generated by weak B field is mutually separated by the Kolmogorov-Arnold-Moser (KAM) tori, those in 3-d systems consist of chaotic zones mutually connected via narrow chaotic channels called the Arnold web (AW) [14-16]. In this case, electrons incoming in an arbitrary direction, which permeate (chaotic) zone by zone in an ergodic way through the AW, can show a global diffusion, i.e., Arnold diffusion (AD). Therefore, a B field increases the transmission channels and the number of stationary points in the integration in Eq.(3), and the resultant integration in Eq.(5) over φ' and φ'' as well as θ' and θ'' indicates $G \propto \frac{e^2}{h} (k_F l)^2$. In general we shall expect a more general result, $G \propto (k_F l)^\beta$ with $1 < \beta \leq 2$, and the exponent β depends on the width of AW. A crucial point is that a B field has increased the exponent in the power-law behavior of the semiclassical conductance in case of the vertical lead orientation. This anomalous phenomenon is just the negative magneto-resistance beyond the weak localization. We here keep to employ the terminology of the *negative magneto – resistance* wherever a B field reduces the zero-field resistance.

On the other hand, for the parallel lead (Q, C) orientation where each electronic trajectory lies almost in the vertical plane, $G \propto \frac{e^2}{h} (k_F l)^2$ is retained, irrespective of the absence or presence of B field; We can expect only the B field-induced increment of the proportionality

constant, which implies a conventional negative magneto-resistance caused by suppression of the weak localization.

Before moving to the next section, we should note: Originally, the negative magneto-resistance was evidenced in the numerical analysis of the conductance of 2-d quantum dots [4]. The accompanying theory, based on the resistance, manipulated backscattering orbits and attributed this phenomenon to the weak localization [4]. Later works [5] based on the conductance, which introduced the small-angle-induced diffraction effect together with an idea of interference between a pair of *partially* time-reversal symmetric orbits [5], swept away a cloud (i.e., breakdown of unitarity) hanging over the above theory. All our results in the present section have been obtained just within the same primitive analysis of the conductance as the original one for 2-d dots.

V. NUMERICAL RESULTS FOR OPEN 3-D ELLIPTIC BILLIARD

To numerically verify the above prediction, we shall choose a SO(2)-symmetric 3-d elliptic shell billiard with the vertical lead orientation. This ellipsoid is a typical example of completely integrable SO(2)-symmetric 3-d billiards. The system in the presence of a symmetry-breaking field B parallel to X axis will be examined. To understand the classical dynamics of electrons, we analyze as a Poincaré surface of section the $X - Z$ plane with longitude S and velocity component v_Z/v^0 as Birkhoff coordinates [17]. ($v^0 = \frac{\hbar\pi}{lm_e}$ is the velocity unit corresponding to the lowest mode.) For $B = 0$, the trajectory from the entrance A with an arbitrary initial velocity vector lying out of $Z - Y$ plane proves to be confined to a torus (see Fig.2(a)), failing to reach the exit C , i.e., $S = \pi/2$. Let $\eta (\equiv \mathfrak{R}/\mathfrak{R}_c)$ be the ratio between the cyclotron radius $\mathfrak{R}_c$ and the characteristic length $\mathfrak{R}$ of the cross-sectional ellipse (see the caption of Fig.1). Then B field can be expressed as $B = \eta B_0$ with $B_0 = m_e v^0 / e \mathfrak{R}$ (B_0 corresponds to $\mathfrak{R}$). In case of a weak field ($B = 0.02B_0$), Poincaré sections for the




FIG. 2. Poincaré sections ($X - Z$ plane) for the trajectories emanating from point A . Longitude S ($-\pi/2 \leq S \leq \pi/2$) and velocity component v_Z/v^0 ($v^0 = \frac{\hbar}{m_e}$) are chosen as Birkhoff coordinates. $S = 0$ and $S = \pi/2$ imply entrance A and exit C , respectively. (a) $B = 0$ and initial velocity $v_X/v^0 = 0.821, v_Y/v^0 = 0.549, v_Z/v^0 = 0.157$; (b),(c) $B = 0.02B_0$ and the same initial velocity as (a): (b) $P = 500$; (c) $P = 17000$. (d)-(f) $B = 0.02B_0$ and initial velocity $v_X/v^0 = 0.843, v_Y/v^0 = 0.538, v_Z/v^0 = 0$: (d) $P = 500$; (e) $P = 4100$; (f) $P = 16000$. P is the number of bouncings at the outer billiard intersecting $X - Z$ plane. For the meaning of B_0 , see the text.

trajectory with the same initial velocity vector as in Fig.2(a) are given in Figs.2(b) and (c)

for increasing numbers of bouncings at the outer billiard wall, and similarly those for the trajectory with the initial velocity vector lying in the $X - Y$ plane are given in Figs.2(d)-(f). One can find: as the B field is switched on, the orbit trajectory first leaves the initial torus, enters an outer layer, and, repeating a similar process, diffuses over more and more distant layers, exhibiting a phenomenon of the Arnold diffusion (AD). Even though a symmetry-breaking field is weak enough to keep the orbit almost straight, AD makes it possible for the trajectories with initial angular momenta $L_Z \neq 0$ to reach the exit C , i.e., $S = \pi/2$ (see Figs.2(c) and (f)) and eventually to contribute to conductance.




FIG. 3. Numerical results for semiclassical conductance versus Fermi wavenumber for A, C leads orientation. Circle and square symbols are for $B = 0.02B_0$ and $B = 0$, respectively. Solid quadratic and dashed linear lines are the corresponding fitted curves for their means.

To proceed to the numerical calculation of semiclassical transmission amplitudes in Eqs.(3,4) for the 3-d elliptic billiard with the vertical lead orientation, let us define the 4×4 monodromy matrix $\mathbf{M}$ as $\begin{pmatrix} \delta \mathbf{r}_\perp'' \\ \delta \mathbf{p}_\perp'' \end{pmatrix} = \mathbf{M} \begin{pmatrix} \delta \mathbf{r}_\perp' \\ \delta \mathbf{p}_\perp' \end{pmatrix}$. At the entrance we introduce a pair of mutually orthogonal coordinates $r_\perp^{1'}$ and $r_\perp^{2'}$ lying in the transverse plane perpendicular to the initial direction of each orbit. Similarly $r_\perp^{1''}$ and $r_\perp^{2''}$ are chosen at the exit. Using elements (M_{ij}) of $\mathbf{M}$ matrix, we rewrite the necessary factors for the numerical computation. For example, $\frac{\partial \theta''}{\partial x''} = \frac{\partial(\theta'', y'')}{\partial(r_\perp^{1'}, r_\perp^{2'})} \frac{\partial(r_\perp^{1'}, r_\perp^{2'})}{\partial(r_\perp^{1''}, r_\perp^{2''})} \frac{\partial(r_\perp^{1''}, r_\perp^{2''})}{\partial(x'', y'')} = \frac{1}{p_F} [M_{41}(\cos \varphi'' \cos \theta'' M_{12} - \sin \varphi'' M_{22}) - M_{42}(\cos \varphi'' \cos \theta'' M_{11} - \sin \varphi'' M_{21})] / (M_{11} M_{22} - M_{12} M_{21})$ and $\frac{\partial \theta'}{\partial x''} = \frac{\partial(\theta', y'')}{\partial(r_\perp^{1''}, r_\perp^{2''})} \frac{\partial(r_\perp^{1''}, r_\perp^{2''})}{\partial(x'', y'')} = \frac{\cos \theta''}{p_F} (\sec \varphi'' \csc \theta'' / M_{14} + \csc \varphi'' / M_{24})$. While $\mathbf{M}$ and ν (Maslov index) can be computed by the method suggested in Refs.[9,18], the first and last rotation angles in $\mathbf{M}$ should be determined by using $r_\perp^{1'}$, $r_\perp^{2'}$ and $r_\perp^{1''}$, $r_\perp^{2''}$ chosen as above, since, after folding the orbit, the initial and final coordinates should become identical. Combining all these factors with Eqs.(3,4), the semiclassical conductance is calculated explicitly. The result, which includes both the classical contribution and the quantum correction in an inseparable way, is plotted in Fig.3. Although the data show small fluctuations, the fitted curves for their means are described by $G \propto \frac{e^2}{h} (k_F l)^2$ and $G \propto \frac{e^2}{h} k_F l$ in cases of $B = 0.02 B_0$ and $B = 0$, respectively, confirming the analytic issue suggested in the previous section.

The increase of the exponent in the k_F dependence of G upon switching on a weak field, which cannot be seen in 2-d cases, provides a numerical evidence that the negative magnetoresistance (: a difference between two lines in Fig.3) in 3-d systems comes from AD as well as the weak-localization correction.

VI. SUMMARY AND DISCUSSIONS

We have investigated semiclassical quantum transport in three-dimensional (3-d) ballistic quantum billiards, and explored novel phenomena that cannot be seen in 2-d billiards. For partially or completely broken-ergodic 3-d billiards such as SO(2) symmetric billiards, the

dependence of the conductance G on the Fermi wavenumber k_F is dramatically changed by the lead orientation. In case of the vertical orientation, a symmetry-breaking weak magnetic field turns out to increase the exponent in k_F dependence of G . In marked contrast to the 2-d case, the anomalous increase of the conductance should be caused by the Arnold diffusion (AD) as well as a weak localization correction.

We should note the following: (i) In an extremely weak field case when the width of Arnold web (AW) is thin enough, we shall see $G \propto (k_F l)^\beta$ with $1 < \beta < 2$. Our numerical calculation in the previous section suggests that, to have $\beta = 2$ in a very large k_F regime, $\mathfrak{R}/\mathfrak{R}_c$ should be no less than 10^{-2} ; (ii) role of a stopper is also essential. Reflection at an inner convex wall should increase the instability of orbits, which is favourable for a genesis of AD; (iii) to clearly observe AD, the symmetry-breaking field should destroy all constants of motion except for the energy. From this viewpoint, it is more advantageous to choose a SO(2)-symmetric 3-d chaotic shell billiard generated from a fully-chaotic 2-d billiard with a hollow by its rotation around the Z axis. The billiards of this kind, which are partially broken-ergodic, are experimentally more accessible than a (completely broken-ergodic) elliptic shell billiard, and all conclusions for the latter should hold for the former as well.

Finally, we point out the role of temperature in observing the effect of AD on the conductance. At finite temperatures, the conductance is related to the scattering matrix as $G = \frac{2e^2}{h} \int_{-\infty}^{\infty} d\epsilon [-\frac{df}{d\epsilon}] \sum_{\mathbf{n}, \mathbf{m}} |t_{\mathbf{n}, \mathbf{m}}|^2$ with f the Fermi distribution function. Then the integration over energy will give a temperature factor $\frac{\pi k_B T L / (\hbar v_F)}{\sinh(\pi k_B T L / (\hbar v_F))}$ with L the length of trajectories. For the vertical lead orientation, this factor suppresses very long trajectories reaching the outgoing lead along the Arnold web, and thereby the negative magneto-resistance is described only by the weak-localization correction. On the other hand, if the temperature is lowered, the self-averaging over k_F is not permissible, leading to a suppression of the weak-localization correction. Then the increment of the magneto-resistance is mainly caused by the effect corresponding to the Arnold diffusion. We hope that a future experiment on 3-d ballistic cavities can verify our prediction.

REFERENCES

- [1] . S. Krylov, *Works on the Foundation of Statistical Physics* (Princeton University Press, Princeton , 1979); Ya. G. Sinai, Russ. Math. Surv. **25**, 137 (1970); P. Gaspard and G. Nicolis, Phys. Rev. Lett. **65**, 1693 (1990).
- [2] . M. Marcus, A. J. Rimberg, R. M. Westervelt, P. F. Hopkins and A. C. Gossard, Phys. Rev. Lett. **69**, 506 (1992); A. M. Chang, H. U. Baranger, L. N. Pfeiffer, and K. W. West, Phys. Rev. Lett. **73**, 2111 (1994).
- [3] special issue on *Chaos and Quantum Transport*: Chaos, Solitons and Fractals **8**, No. **7** and **8** (1997).
- [4] . A. Jalabert, H. U. Baranger, and A. D. Stone, Phys. Rev. Lett. **65**, 2442 (1990). H. U. Baranger, R. A. Jalabert, and A. D. Stone, Phys. Rev. Lett. **70**, 3876 (1993); Chaos. **3**, 665 (1993).
- [5] . L. Aleiner, and A. I. Larkin, Chaos, Solitons & Fractals **8**, 1179 (1997); Y. Takane, and K. Nakamura, J. Phys. Soc. Jpn. **66**, 2977 (1997)).
- [6] . A. Lin, and R. V. Jensen, Phys. Rev. B **53**, 3638 (1996).
- [7] . A. Lin, Chaos, Solitons & Fractals **8**, 995 (1997).
- [8] . P. Bird, R. Akis, D. K. Ferry, D. Vasileska, J. Cooper, Y. Aoyagi, and T. Sugano Phys. Rev. Lett. **82**, 4691 (1999).
- [9] . Primack, and U. Smilansky, Phys. Rev. Lett. **74**, 4831 (1995); Phys. Rep. **327**, 1 (2000).
- [10] . S. Fisher, and P. A. Lee, Phys. Rev. B **23**, 6851 (1981).
- [11] . D. Stone, and A. Szafer, IBM J. Res. Dev. **32**, 384 (1988).
- [12] . C. Gutzwiller, *Chaos in Classical and Quantum Mechanics* (Springer, Berlin, 1990).

- [13] . M. Strutinsky, and A. G. Magner, Sov. J. Part. Nucl. **7**, 138 (1976).
- [14] . V. Chirikov, Phys. Rep. **52**, 263 (1979).
- [15] . J. Lichtenberg, and M. A. Lieberman, *Regular and Stochastic Motion* (Springer, New York, 1983).
- [16] . M. Zaslavsky, R. Z. Sagdeev, D. A. Usikov, and A. A. Chernikov, *Weak Chaos and Quasi-Regular Patterns* (Cambridge University Press, Cambridge , 1991).
- [17] . D. Birkhoff, *Dynamical Systems* (American Mathematical Society , Providence, RI, 1927).
- [18] . Sieber, Nonlinearity **11**, 1607 (1998).
- [19] . W. J. Beenakker, and H. van Houten, Phys. Rev. Lett. **63**, 1857 (1989).